## **Oregon Status Factors**

Elcode NFSM000027

Gname CLAVARIADELPHUS OCCIDENTALIS

**Gcomname** 

### **Number of Occurrences**

D = 81 - 300

Comments There are 35 sites for Oregon in OHIC database and Methven 1990 examined 17 collections.

So, there are 60 records of this species. Seems to be a fairly common species throughout its

range. Scattered to gregarious, in coniferous or mixed deciduos-coniferous forests.

## **Number of Occurrences with Good Viability**

D = Some (13-40) occurrences with good viability

Comments There are 19 occurrences in protected sites.

## **Population Size**

B = 50-250 individuals

Comments

## **Range Extent**

H = > 2,500,000 km2 (greater than 1,000,000 square miles)

Comments Range is across western Oregon.

## **Area of Occupancy**

B = 0.4-4 km 2 (about 100-1,000 acres)

LB = 4-40 km (about 2.5-25 miles)

Comments

# Long-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Long-term trend in population, range, area occupied, or number or condition of occurrences unknown

Comments

## Short-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Short-term trend in population, range, area occupied, and number and condition of occurrences unknown.

Comments

#### **Threats**

C = Substantial, non-imminent threat. Threat is moderate to severe but not imminent (> 10 years) for most of the population, occurrences, or area.

Scope High Severity High Immediacy Unknown

Comments This is a mycorrhizal species associated with coniferous and deciduous forest. Susceptible to logging activities.

## **Number of Appropriately Protected and Managed Occurrences**

D = Many (13-40) occurrences appropriately protected and managed

Comments There are 19 occurrences in protected sites.

## **Intrinsic Vulnerability**

A = Highly Vulnerable. Species is slow to mature, reproduces infrequently, and/or has low fecundity such that populations are very slow (> 20 years or 5 generations) to recover from decreases in abundance; or species has low dispersal capability such that extirpated populations are unlikely to become reestablished through natural recolonization (unaided by humans). Ecological community occurrences are highly susceptible to changes in composition and structure that rarely if ever are reversed through natural processes even over substantial time periods (> 100 years).

Comments This is a mycorrhizal species susceptible to logging activities.

## **Environmental Specificity**

B = Narrow. Specialist or community with key requirements common.

Comments Mycorrhizal species

## **Other Considerations**

ORNHIC - Not Listed.

Edition 11/12/2002 Edauthor Efren Cazares

Grank Date 11/12/2002

#### **Greasons**

Common mycorrhizal species throughout western OR. 19 out of 43 occurrences are in protected sites. Apparently secure in OR.

#### **BCD Sources**

Castellano, M.A., E. Cazares, B. Fondrick and T. Dreisbach. 2002. Handbook to additional fungal species of special concern in the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-xxx. Portland, OR; U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. xxx p. (In press)& Methven, A.S. 1990. The genus Clavariadelphus in North America. Bibliotheca Mycologica. J. Cramer. pp 192.

#### **New Sources**