

Conservation Status Assessment

Scientific Name: *Gymnopilus punctifolius*

Classification: Fungus

Assessment area: California

Heritage Rank: **S3**

Rank Date: 11/21/2016

Rank Reasons: Several sites, some threats and vulnerability.

Range Extent: E = 5,000-20,000 sq km (~2,000-8,000 sq mi)

Comments: Range extent is 14,267 sq. Km. This area is restricted to the west slopes of the northern California coast range with most sites in redwood forests.

Population Size: Not assessed

Comments: None

Number of Occurrences: B = 6 - 20

Comments: 12 occurrences in California

Area of Occupancy: D = 6-25 4-km² grid cells

Comments: Occupies 19 grid squares in California

Good Viability: C = Few (4-12) occurrences with excellent or good viability or ecological integrity

Comments: Most sites in the permanently protected California Redwoods State Parks, but only 12 sites overall.

Environmental Sensitivity: BC = Narrow to moderate

Comments: As per L. L. Norvell in 2002 ranking - Saprophytic on cubical rotted coniferous wood, debris and rich humus (Hesler 1969; Castellano et al 1999) in mid-successional (Norvell pers. comm. 2002) to LSOG forests where coarse woody debris is preserved under closed canopies. Its precise biological and ecological requirements are unknown. One collection from Oregon made from a 55 year old forest came from a well-protected and very moist rotting log in an area that was clear-cut the next year before the fall mushroom season. It has not been observed in the ensuing four years. (Norvell & Exeter 2003).

Short Term Trends: Not Evaluated

Comments: None

Long Term Trends: Not Evaluated

Comments: None

Threat Impact: D = Low

Comments:

Known sites in California are mostly in Redwood State Parks, which protects them from logging. The redwood belt may be vulnerable to climate change impacts.

Intrinsic Vulnerability: B = Moderately vulnerable

Comments: According to the 2002 ranking by L. L. Norvwell, GYPU2 is inferred to be particularly vulnerable to removal of its substrate, clean cutting with burning or removal of all coarse woody debris, or logging activities that appreciably open the canopy and expose the substrate to sun and wind. It is also vulnerable to alteration of microhabitats and microclimate regimes (stream diversion, road construction, development).

Calculated Rank: S3

Rank Author: Michael Russell

Rank Reviewer: Michael Russell

References:

No additional references listed.

Definitions and Resources:

Rank Prefixes

G Global rank, applied to taxon's full geographic range

S State rank, applied to taxon's range within the designated state

Rank Values

1 Critically imperiled

2 Imperiled

3 Vulnerable

4 Apparently secure, uncommon but not rare

5 Secure, common, abundant, and widespread

Suggested citation:

Oregon Biodiversity Information Center. 2017. Fungi Conservation Status Assessments. Institute for Natural Resources, Portland State University and Oregon State University. Portland, Oregon and Corvallis, Oregon.

More assessments available at <http://inr.oregonstate.edu/orbic/rare-species/ranking-documentation>

Element rank calculator resources at <http://www.natureserve.org/conservation-tools/conservation-rank-calculator>

Oregon Biodiversity Information Center, Institute for Natural Resources

Oregon State University and Portland State University

Mail Stop: INR, P.O. Box 751

Portland, OR 97207-0751

(503)-725-9950

<http://inr.oregonstate.edu/orbic>

<http://inr.oregonstate.edu/>