

# Heritage Ranking Form - Global Rank

**Scientific Name:** Phaeocollybia dissiliens

**Common Name:** Fungus

**Classification:** Fungus

**Range Extent:** F = 20,000-200,000 sq km (~8,000-80,000 sq mi)

Comments: about 10,000 sq. mi., mainly along the Pacific Coast and Coast Range from Cape Lookout, Tillamook County, Oregon, south to Del Norte County, California. A few occurrences in western Oregon Cascades.

**Area of Occupancy:** E = 26-125 4-km<sup>2</sup> grid cells

Comments: 27 grid cells

**Number of Occurrences:** C = 21 - 80

Comments: About 28 occurrences

**Population Size:** U = Unknown

Comments: None

**Good Viability:** U = Unknown

Comments: None

**Environmental Specificity:** B = Narrow. Specialist or community with key requirements common

Comments: Mycorrhizal in moist coniferous forests of Picea, Tsuga, Pseudotsuga and Abies (Norvell & Exeter 2008)

**Short Term Trends:** U = Unknown

Comments: None

**Long Term Trends:** U = Unknown

Comments: None

**Threat Impact:** C = Medium

Comments: Species grow in coniferous forests of the western US which has been managed primarily for timber production.

**Intrinsic Vulnerability:** Not Evaluated

Comments: None

**Heritage Rank:** G3

Comments: Currently known from 28 occurrences, distributed from NW Oregon to northern California, mainly along the coast or in the Coast Range.

Rank Notes: None

Reference: Norvell, L.L. and R.L. Exeter. 2008. Phaeocollybia of Pacific Northwest North America. USDI BLM/OR/WA/GI-08/100-1792, Salem, Oregon 228 p.

Rank Date: 6/15/2010

Rank Author: Vrilakas, Sue

Oregon Biodiversity Information Center  
Institute for Natural Resources  
Portland State University  
Mail Stop: INR, PO Box 751  
Portland, OR 97207-0751  
503-725-9950  
[www.orbic.pdx.edu](http://www.orbic.pdx.edu)