

# Oregon Status Factors

**Elcode** NFSM000137

**Gname** RAMARIA AMYLOIDEA

**Gcomname**

## Number of Occurrences

B = 6 - 20

**Comments** 6 sites in the Oregon Cascades.

## Number of Occurrences with Good Viability

C = Few (4-12) occurrences with good viability

**Comments** There are 6 occurrences in protected sites. The rest of the sites are subject to logging activities that could affect the populations.

## Population Size

A = 1-50 individuals

**Comments**

## Range Extent

E = 5,000-20,000 km<sup>2</sup> (about 2,000-8,000 square miles)

**Comments** Occurs in the Oregon Cascades.

## Area of Occupancy

A = <0.4 km<sup>2</sup> (less than about 100 acres)

LA = <4 km (less than about 2.5 miles)

**Comments** Associated to *Abies* spp., *Pseudotsuga menziesii* and *Tsuga heterophylla*.

## Long-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Long-term trend in population, range, area occupied, or number or condition of occurrences unknown

**Comments**

## Short-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Short-term trend in population, range, area occupied, and number and condition of occurrences unknown.

**Comments**

## Threats

C = Substantial, non-imminent threat. Threat is moderate to severe but not imminent (> 10 years) for most of the population, occurrences, or area.

Scope High Severity High Immediacy Unknown

**Comments** This species is strictly associated to its hosts (mycorrhizal). Tree removal and compaction (Logging activities) will depauperate the populations. This species is known to occur in late successional forest only.

## Number of Appropriately Protected and Managed Occurrences

C = Several (4-12) occurrences appropriately protected and managed

**Comments** There are 6 occurrences in protected sites. The rest of the sites are subject to logging activities that could affect the populations.

## Intrinsic Vulnerability

A = Highly Vulnerable. Species is slow to mature, reproduces infrequently, and/or has low fecundity such that populations are very slow (> 20 years or 5 generations) to recover from decreases in abundance; or species has low dispersal capability such that extirpated populations are unlikely to become reestablished through natural recolonization (unaided by humans). Ecological community occurrences are highly susceptible to changes in composition and structure that rarely if ever are reversed through natural processes even over substantial time periods (> 100 years).

**Comments** This species is strictly associated to its hosts (mycorrhizal). Tree removal and compaction (Logging activities) will depauperate the populations. This species is known to occur in late successional forest only.

## Environmental Specificity

B = Narrow. Specialist or community with key requirements common.

**Comments** Mycorrhizal species

## Other Considerations

ORNHIC List 2

**Edition** 11/5/2002 **Edauthor** Efren Cazares

**Grank** S2? **Grank Date** 11/5/2002

## Reasons

Only found in 6 locations in the Oregon Cascades and a mycorrhizal species that depends on late successional forest.

## BCD Sources

Castellano, M.A., E. Cazares, B. Fondrick and T. Dreisbach. 2002. Handbook to additional fungal species of special concern in the Northwest Forest Plan. Gen. Tech. Rep. PNW-GTR-xxx. Portland, OR; U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. xxx p. (In press) and Petersen, R.H. 1981. Ramaria subgenus Echinoramaria. J. Cramer. Pp. 261.

## New Sources