Washington Status Factors

Elcode NFSM000148

Gname RAMARIA GELATINIAURANTIA

Gcomname

Number of Occurrences

A = 1 - 5

Comments Known from 5 locations in WA.

Number of Occurrences with Good Viability

B = Very few (1-3) occurrences with good viability

Comments Mycorrhizal species and susceptible to logging activities. There are two locations in WA National

Parks that might be protected.

Population Size

A = 1-50 individuals

Comments

Range Extent

F = 20,000-200,000 km 2 (about 8,000-80,000 square miles)

Comments Known from 5 locations. From Olympic peninsula to Gifford Pinchot National Forests.

Area of Occupancy

A = <0.4 km 2 (less than about 100 acres)

LA = <4 km (less than about 2.5 miles)

Comments

Long-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Long-term trend in population, range, area occupied, or number or condition of occurrences unknown

Comments

Short-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Short-term trend in population, range, area occupied, and number and condition of occurrences unknown.

Comments

Threats

C = Substantial, non-imminent threat. Threat is moderate to severe but not imminent (> 10 years) for most of the population, occurrences, or area.

Scope High Severity High Immediacy Unknown

Comments Mycorrhizal species associated to late successional forest of Douglas fir and western Hemlock susceptible to logging activities.

Number of Appropriately Protected and Managed Occurrences

B = Few (1-3) occurrences appropriately protected and managed

Comments Populations in Olympic and Mt. Rainier National Parks are protected.

Intrinsic Vulnerability

A = Highly Vulnerable. Species is slow to mature, reproduces infrequently, and/or has low fecundity such that populations are very slow (> 20 years or 5 generations) to recover from decreases in abundance; or species has low dispersal capability such that extirpated populations are unlikely to become reestablished through natural recolonization (unaided by humans). Ecological community occurrences are highly susceptible to changes in composition and structure that rarely if ever are reversed through natural processes even over substantial time periods (> 100 years).

Comments Mycorrhizal species susceptible to logging activities.

Environmental Specificity

B = Narrow. Specialist or community with key requirements common.

Comments Mycorrhizal species

Other Considerations

Edition 11/7/2002 Edauthor Efren Cazares

Grank S3 Grank Date 11/7/2002

Greasons

Wide range but not a common mycorrhizal species vulnerable to logging activities. Only 2 out of 5 sites are protected. More studies are needed to determine its rarity or abundance in WA.

BCD Sources

Castellano, M.A., J.E. Smith, T. O'Dell, E. Cazares and S. Nugent. 1999. Handbook to Strategy 1 Fungal species in the Northwest Forest Plan. USDA, Forest Service Pacific Northwest Research Station, Portland, OR. GTR PNW-GTR-476. & Marr, C.D. and Stuntz, D.E. 1973. Ramaria of Western Washington. Biblio. Mycol. 38:1-232.

New Sources