

	<b>Species:</b>	<b>Index Result:</b>
Scientific Name	<i><b>Brodiaea terrestris</b></i>	<b>Extremely Vulnerable</b>
Common Name	<b>Dwarf brodiaea</b>	<b>Confidence Very High</b>
Taxonomic Group	Vascular Plant	(based on entered data)
Geographic Area	Western (coastal) Oregon	Date Assessed 5/6/2020
		GRank G4G5
Cave/Ground Water Obligate: No		SRank S2
Migratory area included in assessment: No		Assessor Sue Vrillakas

**Climate Change Vulnerability Index Values:** (greatest score shown when range was selected)

Category	Factor	Score	Comments
Temperature Scope (predicted increase)	A >6.0F	0	
	A 5.5F	0	
	A 5.1F	0	
	A 4.5F	0	
	A 3.9F	0	
	A <3.9F	100	
Hamon AET:PET Moisture Metric Scope	< -0.119	0	
	-0.119	0	
	-0.096	100	
	-0.073	0	
	-0.05	0	
	>-0.028	0	
Sea level rise	B1	GI	Grows on coastal sand dunes, open areas, lowlands and sandy bluffs; stronger storm surges and frequency may change landscape
Natural barriers	B2a	Inc	Limited to coastal area but maybe able to shift its range
Anthropogenic barriers	B2b	N	
Climate Change mitigation	B3	U	
Dispersal/Movement	C1	Inc	Found in region where temperature variation historically small  Highest value: 80.9369; lowest 61.1821; difference=19.7548  Restricted to sand substrate
Historical thermal niche	C2ai	GI	
Physiological thermal niche	C2aai	N	
Historical hydrological niche	C2bi	SI	
Physiol. hydrological niche	C2bii	N	
Disturbance dependence	C2c	U	
Ice/snow dependence	C2d	N	
Physical habitat restrictions	C3	SI	
Other spp create habitat	C4a	N	
Dietary Versatility	C4b	U	
Pollinator Versatility	C4c	U	
Other spp for dispersal	C4d	N	
Pathogen sensitivity	C4e	N	
Competition sensitivity	C4f	N	
Interspecific Relationship	C4g	U	
Measured genetic variation	C5a	U	
Bottlenecks	C5b	U	
Plant reproductive system	C5c	U	
Phenological response	C6	U	

Documented response	D1	U	
Modeled change	D2	U	
Modeled overlap	D3	U	
Modeled protected areas	D4	U	

**Additional Notes:**

Range map created using ArcMap Minimum Mapping Boundary-Convex Hull on ORBIC element occurrence data, 4-29-2020 export, then edited to match more closely the coast line. Climate and precipitation data from Climate Wizard using the A1B emissions scenario and ensemble average general circulation model: Historical = 1951-2006; Future = mid-century (2050s); Hamon AET:PET moisture metric (Hamon 1961).

**References:****Data sources and notes:**

Range map created using ArcMap Minimum Mapping Boundary-Convex Hull on ORBIC element occurrence data. Climate and precipitation data from Climate Wizard using the A1B emissions scenario and ensemble average general circulation model: Historical = 1951-2006; Future = mid-century (2050s); Hamon AET:PET moisture metric (Hamon 1961).

Detailed definitions of criteria and methodology can be found in the documentation at <http://www.natureserve.org/conservation-tools/climate-change-vulnerability-index>

**Legend and Definitions:**

<b>Affect to Vulnerability:</b>
GI = Greatly increase
Inc = Increase
SI = Somewhat increase
N = Neutral
U = Unknown

**Index Scores:**

<b>Extremely Vulnerable:</b> Abundance and/or range extent within geographical area assessed extremely likely to substantially decrease or disappear by 2050.
<b>Highly Vulnerable:</b> Abundance and/or range extent within geographical area assessed likely to decrease significantly by 2050.
<b>Moderately Vulnerable:</b> Abundance and/or range extent within geographical area assessed likely to decrease by 2050.
<b>Less Vulnerable:</b> Available evidence does not suggest that abundance and/or range extent within the geographical area assessed will change (increase/decrease) substantially by 2050. Actual range boundaries may change.
<b>Insufficient Evidence:</b> Information entered about a species' vulnerability is inadequate to calculate an Index score.

**Citation:**

Oregon Biodiversity Information Center. 2020. Climate Change Vulnerability Index assessment for Dwarf brodiaea (*Brodiaea terrestris*). Institute for Natural Resources, Portland State University, Portland, OR.