

## Oregon Status Factors

**Elcode** NLCAL00008  
**Gname** CHAENOTHECA SUBROSCIDA  
**Gcomname** lemondrop whiskers

### Number of Occurrences

**Comments** Number of known occurrences in Oregon = 7. All are recent and are mostly extant.

### Number of Occurrences with Good Viability

U = Unknown what number of occurrences with good viability

**Comments**

### Population Size

U = Unknown

**Comments**

### Range Extent

H = > 2,500,000 km<sup>2</sup> (greater than 1,000,000 square miles)

**Comments** Restricted to the range of the Northern Spotted Owl in Oregon.

### Area of Occupancy

A = <0.4 km<sup>2</sup> (less than about 100 acres)

LA = <4 km (less than about 2.5 miles)

**Comments** Occupancy for epiphytic lichens and fungi can be difficult to estimate, particularly for calicioid species (including this species) which often occur as colonies covering only a few square centimeters on single tree trunk within a stand and then again several hundred meters away. The occupancy given above is roughly estimated as the total worldwide distribution of the species; the actual coverage of the species condensed so as to be continuous may not be much more than 1 hectare.

### Long-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

B = Large Decline (decline of 75-90%)

**Comments** Most calicioid lichens and fungi inhabit aged bark or wood in sheltered locations protected from direct rain interception. This species is very restricted to the bark of old trees; in the Pacific Northwest of North America, most known occurrences are on conifers > 200 years old, with occasional occurrences on younger trees (e.g. 150 years old) southward in the Klamath region (Peterson unpublished data). Removal of old forests in North America and through the rest of the species' distribution has undoubtedly had severe impacts on the number of populations, population sizes, and average dispersal distance necessary to colonize new substrates.

## Short-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

D = Declining. Decline of 10-30% in population, range, area occupied, and/or number or condition of occurrences

**Comments** With advances in conservation, the removal of old-growth forests throughout the species range is slowing, but has not stopped.

## Threats

B = Moderate and imminent threat. Threat is moderate to severe and imminent for a significant proportion (20-60%) of the population, occurrences, or area. Ecological community occurrences are directly impacted over a moderate area, either causing irreversible damage or requiring a long-term recovery.

**Scope** Moderate                      **Severity** High                      **Immediacy** High

**Comments** Worldwide, the species has gone through drastic declines since pre-industrial times. The Pacific Northwest, due to logging, has been no exception. However, the rate of loss in the Pacific Northwest has slowed. Although little is known about the reproductive and dispersal biology of this species, it is thought that the species can overcome some habitat fragmentation. Since the species is quite rare in the Pacific Northwest, and apparently in other parts its distribution, the species is at some risk of extirpation or extinction. That, combined with its strong association with old growth warrens significant consideration in conservation actions.

## Number of Appropriately Protected and Managed Occurrences

U = Unknown whether any occurrences are appropriately protected and managed

**Comments**

## Intrinsic Vulnerability

A = Highly Vulnerable. Species is slow to mature, reproduces infrequently, and/or has low fecundity such that populations are very slow (> 20 years or 5 generations) to recover from decreases in abundance; or species has low dispersal capability such that extirpated populations are unlikely to become reestablished through natural recolonization (unaided by humans). Ecological community occurrences are highly susceptible to changes in composition and structure that rarely if ever are reversed through natural processes even over substantial time periods (> 100 years).

**Comments** Given high vulnerability rank because it will not return to a forest for a century or more after a stand-initiating disturbance. Although the species is limited to substrates that are very slow to develop and the maturation time required between colonization and reproduction is unknown, the species does demonstrate a remarkable ability to disperse to appropriate substrates once they are available, even when those substrates are rather isolated. This may be due to use of a dispersal vector such as birds or arthropods which target similar habitats.

## Environmental Specificity

A = Very Narrow. Specialist or community with key requirements scarce.

**Comments**

## Other Considerations

ORNHIC - List 3.

**Edition** 2/20/2003                      **Edauthor** Eric B. Peterson

**Grank** S2S3                      **Grank Date** 11/22/2002

## Greasons

Known from 7 occurrences, all in old-growth forests, and only one in an LSR in Oregon. The other 6 are currently protected, have been excluded from timber sales, but with the change in status, they may not be secure. If these known locations become at risk, the status will become S2.

## BCD Sources

### New Sources

ISMS (Interagency Species Management System). 5 August, 2002.

Peterson, E. B. (Search of personal herbarium on 1 November, 2002) Address: Nevada Natural Heritage Program, 1550 E. College Parkway, Carson City, NV

Rikkinen, J. 2003. Calicioid lichens and fungi in the forests and woodlands of western Oregon. *Annales Botanici Fennici* (accepted, should come out in the first volume of 2003).

Selva, S., 7 November 2002. Personal communication. Address: Division of Natural and Behavioral Sciences, University of Maine at Fort Kent, Fort Kent, ME.

Tibell, L. 1975. The Caliciales of boreal North America. *Symbolae Botanicae Upsalienses* 21(2): 1-128.

Tibell, L. 1999. Caliciales. *Nordic Lichen Flora* 1: 20-93.