

Conservation Status Assessment

Scientific Name: *Gymnopilus punctifolius*

Classification: Fungus

Assessment area: Oregon

Heritage Rank: **S3**

Rank Date: 11/21/2016

Rank Reasons: Several sites but has vulnerabilities.

Range Extent: F = 20,000-200,000 sq km (~8,000-80,000 sq mi)

Comments: Range is 47070 sq. Km. There are known sites just across the Idaho border so the actual range may be larger, but not enough to cause a different score

Population Size: Not assessed

Comments: None

Number of Occurrences: C = 21 - 80

Comments: 50 occurrences in Oregon

Area of Occupancy: E = 26-125 4-km² grid cells

Comments: 59 occupied grid squares in Oregon

Good Viability: B = Very few (1-3) occurrences with excellent or good viability or ecological integrity

Comments: Oregon sites are in BLM and forest service lands, but none are in any permanently protected areas.

Environmental Sensitivity: BC = Narrow to moderate

Comments: As per L. L. Norvell in 2002 ranking - Saprophytic on cubical rotted coniferous wood, debris and rich humus (Hesler 1969; Castellano et al 1999) in mid-successional (Norvell pers. comm. 2002) to LSOG forests where coarse woody debris is preserved under closed canopies. Its precise biological and ecological requirements are unknown. One collection from Oregon made from a 55 year old forest came from a well-protected and very moist rotting log in an area that was clear-cut the next year before the fall mushroom season. It has not been observed in the ensuing four years. (Norvell & Exeter 2003).

Short Term Trends: Not Evaluated

Comments: None

Long Term Trends: Not Evaluated

Comments: None

Threat Impact: C = Medium

Comments:

Threats include timber harvest which could impact the over 80% of sites that are not in national parks on as short as a 30 - 40 year rotation, fire which if it occurs on the historic fire return interval may occur between every 25 to over 300 years depending on the site, and climate change which may increase the fire return interval or dry out the forest at an unknown number of sites.

Intrinsic Vulnerability: B = Moderately vulnerable

Comments: According to the 2002 ranking by L. L. Norvwell, GYPU2 is inferred to be particularly vulnerable to removal of its substrate, clean cutting with burning or removal of all coarse woody debris, or logging activities that appreciably open the canopy and expose the substrate to sun and wind. It is also vulnerable to alteration of microhabitats and microclimate regimes (stream diversion, road construction, development).

Calculated Rank: S3

Rank Author: Michael Russell

Rank Reviewer: Michael Russell

References:

No additional references listed.

Definitions and Resources:

Rank Prefixes	
G	Global rank, applied to taxon's full geographic range
S	State rank, applied to taxon's range within the designated state
Rank Values	
1	Critically imperiled
2	Imperiled
3	Vulnerable
4	Apparently secure, uncommon but not rare
5	Secure, common, abundant, and widespread

Suggested citation:

Oregon Biodiversity Information Center. 2017. Fungi Conservation Status Assessments. Institute for Natural Resources, Portland State University and Oregon State University. Portland, Oregon and Corvallis, Oregon.

More assessments available at <http://inr.oregonstate.edu/orbic/rare-species/ranking-documentation>

Element rank calculator resources at <http://www.natureserve.org/conservation-tools/conservation-rank-calculator>

Oregon Biodiversity Information Center, Institute for Natural Resources
 Oregon State University and Portland State University
 Mail Stop: INR, P.O. Box 751
 Portland, OR 97207-0751 <http://inr.oregonstate.edu/orbic>
 (503)-725-9950 <http://inr.oregonstate.edu/>