

Oregon Status Factors

Elcode NFSM000123
Gname PHAEOCOLLYBIA OLIVACEA
Gcomname

Number of Occurrences

C = 21- 80

Comments There are ~53 total known occurrences of PHOL in Oregon, of which at least 49 are believed extant (Norvell 1998ac, 2002; ISMS 2002 database & GIS map for PHOL).

Number of Occurrences with Good Viability

E = Many (41-125) occurrences with good viability

Comments There are 49 known extant occurrences.

Population Size

U = Unknown

Comments Records reflect only species occurrence, i.e. fruitbodies, not numbers of individuals. Genets of ectomycorrhizal fungi cannot be delimited without DNA sampling.

Range Extent

F = 20,000-200,000 km² (about 8,000-80,000 square miles)

Comments Endemic to the northern spotted region in western North America, in Oregon PHOL ranges from the Pacific coast (from 45° latitude N south to the California border) east to the lower west slope of the Cascade Mountains. (Norvell 1998ac, ISMS 2002 database and GIS map for PHOL).

Area of Occupancy

U = Unknown

LU = Unknown

Comments Occupancy is highly spotty and cannot be extrapolated for this organism, which appears restricted to fairly complex environments. For fungi can only estimate area occupancy from fruitbodies as vegetative organism is underground had has unknown biological and ecological requirements that determine how and when ectomycorrhizal associations are formed with coniferous host trees. There are large areas of unsuitable habitat within the overall range.

Long-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

D = Moderate Decline (decline of 25-50%)

E = Relatively Stable ($\pm 25\%$ change)

Comments Due to the spotty nature of the occurrences, it is difficult to project a long-term trend in population size, extent of occurrence, or the area of occupancy. Ectomycorrhizal fungal stability depends on the stability of the coniferous partners so that the fungus and forest trends coincide. PHOL occurs most frequently in mixed fagaceous/coniferous forests, apparently forming preferential symbiotic associations with *Quercus* or *Lithocarpus* spp. Because of this inferred preference, PHOL

occurrences may be greatly affected by the advent of sudden oak death *Phytophthora* in southern Oregon over the long term. Current populations appear healthy and are inferred to be long-lived; individuals are less dependent upon spore dispersal than with mycelial interactions with other individuals and their mycorrhizal partners. (Norvell 1998ab, pers comm 2002). The populations are unusual, but not uncommon.

Short-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

E = Stable. Population, range, area occupied, and/or number or condition of occurrences unchanged or remaining within $\pm 10\%$ fluctuation

Comments Ectomycorrhizal fungal trends are inextricably linked to the trends for the symbiotic partners, in this case trees. Road construction, hot fires, clearcutting, or development that displaces the tree partner will compromise both tree and fungus. Current occurrences of PHOL are relatively common, with at least 16 occurring in protected forest reserves; therefore *P. olivacea* is believed to be stable and possibly expanding over the short term, provided no human interference or hot fires imperil the host forests and they remain free of the sudden oak death *Phytophthora*.

Threats

D = Moderate, non-imminent threat. Threat is moderate to severe but not imminent for a significant portion of the population, occurrences, or area.

Scope Moderate Severity High Immediacy Low

Comments Whatever threatens the extant forest will threaten the fungus. This species is usually found in complex mixed oak-coniferous forests. Populations are inferred to be long-lived from the fact that collections have been made in or near the type locality from 1956 to 2000 (Norvell 1998ac, ISMS 2002). Like the forest, PHOL is threatened by hot fires, road construction or other development, and clearcutting, but appears to be able to withstand light to moderate thinning (Norvell pers comm 2002). The most serious long-term threat is the predicted rapid spread of the "sudden oak death" *Phytophthora* to the oak partner trees in southern Oregon.

Number of Appropriately Protected and Managed Occurrences

Comments ISMS 2002 and Norvell (1998ac) cite approximately 16 occurrences (RANK D) in protected areas: 1 in permanent protected preserves, 10 in late-successional reserves, and ~5 in riparian reserves. If governmental management policies dictate opening late-successional and/or riparian reserves to clearcutting, road construction, or other development, the number of protected and managed occurrences could dramatically decrease to 1 (Rank B). Additionally, many sites in temporary reserves may not be managed appropriately at the present time.

Intrinsic Vulnerability

A = Highly Vulnerable. Species is slow to mature, reproduces infrequently, and/or has low fecundity such that populations are very slow (> 20 years or 5 generations) to recover from decreases in abundance; or species has low dispersal capability such that extirpated populations are unlikely to become reestablished through natural recolonization (unaided by humans). Ecological community occurrences are highly susceptible to changes in composition and structure that rarely if ever are reversed through natural processes even over substantial time periods (> 100 years).

B = Moderately Vulnerable. Species exhibits moderate age of maturity, frequency of reproduction, and/or fecundity such that populations generally tend to recover from decreases in abundance over a period of several years (on the order of 5-20 years or 2-5 generations); or species has moderate dispersal capability such that extirpated populations generally become reestablished through natural recolonization (unaided by humans). Ecological community occurrences may be susceptible to changes in composition and structure but tend to recover through natural processes given reasonable time (10-100 years).

Comments Ectomycorrhizal fungal vulnerability is linked to that of the symbiotic partner (here oak+conifers).

This fungus is thought to be long-lived, but also relatively slow-growing; therefore climax communities occur in older stands. (Norvell 1998ac). It is vulnerable to anything that threatens the forest habitat, including hot fires, road construction and development, clearcutting and the "sudden oak death" Phytophthora in southern Oregon. (Norvell pers comm 2002).

Environmental Specificity

B = Narrow. Specialist or community with key requirements common.

Comments PHOL is generally found in complex mid to late-successional/old growth coniferous rainforests where it forms symbiotic partnerships with *Quercus* or *Lithocarpus* spp and possibly also members of the Pinaceae (it rarely is found in strictly fagaceous or coniferous stands). Its precise biological and ecological requirements are unknown. It generally is found in the more southern part of the northern spotted owl region, and fruits on soil in early to late autumn producing arcs of closely gregarious fruitbodies. It is like all *Phaeocollybias* in its extremely patchy in distribution.. (Norvell 1998ac, Norvell 2002).

Other Considerations

ORNHIC - List 1. There are no known synonyms for the species. This fungus may be highly receptive to the sudden oak death which would attack its inferred preferred mycorrhizal partner. Additional occurrences are to be expected in late-successional or unexplored old-growth forests. There are several currently protected sites.

Edition 11/19/2002 **Edauthor** Lorelei L Norvell

Grank S2 **Grank Date** 11/19/2002

Greasons

PHOL is unusual but not uncommon in southern Oregon, with 49 known extant occurrences, of which 16 lie in currently protected forest reserves. It is endemic to the Pacific Northwest northern spotted owl region. Its patchy distribution precludes estimation of population size and area of occupancy. It usually mixed fagaceous+pinaceous forests, occasionally in pure coniferous stands. Known extant populations appear relatively stable, but the approach of the "sudden oak death" Phytophthora in southern Oregon imperils those in mixed forests. Unprotected occurrences are possibly imperiled also by road construction & development and clearcutting or heavy thinning. Moderate to light thinning is not considered a threat. All occurrences are imperiled by hot fires.

BCD Sources

New Sources

Norvell. 1998a. The biology and taxonomy of Pacific Northwest species of *Phaeocollybia* Heim. 391 pp. ALSO Norvell. 1998b. . Observations on the development, morphology, and biology of *Phaeocollybia*. *Mycological Research* 102:615-630. ALSO Norvell. 1998c. ROD: Strategy 3 Fungal Species Evaluation (11 gilled Basidiomycete Strategy 3 species). Unpubl. report on file at the Regional Mycology Lab, Corvallis, Oregon. ALSO Norvell. 2002. *Phaeocollybia* in western North America 3: *Mycotaxon* 81: 25-112. ALSO ISMS 2002: database + GIS map on PHOL.