

# Oregon Status Factors

**Elcode** NFSM000160  
**Gname** RAMARIA SPINULOSA VAR DIMINUTIVA  
**Gcomname**

## Number of Occurrences

A = 1 - 5

**Comments** There is only 1 record in OR.

## Number of Occurrences with Good Viability

U = Unknown what number of occurrences with good viability

**Comments** This is a mycorrhizal species associated to late successional forests. Unknow forest management activities.

## Population Size

A = 1-50 individuals

**Comments**

## Range Extent

D = 1,000-5,000 km<sup>2</sup> (about 400-2,000 square miles)

**Comments** There is only one record from OR. This species is reported as *R. funosiavellana* in Marr and Stuntz 1973.

## Area of Occupancy

A = <0.4 km<sup>2</sup> (less than about 100 acres)

LA = <4 km (less than about 2.5 miles)

**Comments**

## Long-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Long-term trend in population, range, area occupied, or number or condition of occurrences unknown

**Comments**

## Short-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Short-term trend in population, range, area occupied, and number and condition of occurrences unknown.

**Comments**

## Threats

A = Substantial, imminent threat. Threat is moderate to severe and imminent for most (> 60%) of the population, occurrences, or area. Ecological community occurrences are directly impacted over a widespread area, either causing irreversible damage or requiring long term recovery

Scope High

Severity High

Immediacy High

Comments This is mycorrhizal species associated to late successional forests.

## Number of Appropriately Protected and Managed Occurrences

A = None. No occurrences appropriately protected and managed

Comments The sites where it occurs are not protected.

## Intrinsic Vulnerability

A = Highly Vulnerable. Species is slow to mature, reproduces infrequently, and/or has low fecundity such that populations are very slow (> 20 years or 5 generations) to recover from decreases in abundance; or species has low dispersal capability such that extirpated populations are unlikely to become reestablished through natural recolonization (unaided by humans). Ecological community occurrences are highly susceptible to changes in composition and structure that rarely if ever are reversed through natural processes even over substantial time periods (> 100 years).

Comments This is a mycorrhizal species and susceptible to logging activities.

## Environmental Specificity

B = Narrow. Specialist or community with key requirements common.

Comments Mycorrhizal species

## Other Considerations

ORNHIC List 1

**Edition** 11/10/2002

**Edauthor** Efren Cazares

**Grank** S1?

**Grank Date** 11/10/2002

## Reasons

There is only 1 record in OR. Endemic and rare to late successional in the PNW forests. Susceptible to logging activities. More studies are needed to determine its rarity or abundance in the PNW.

## BCD Sources

Castellano, M.A., J.E. Smith, T. O'Dell, E. Cazares and S. Nugent. 1999. Handbook to Strategy 1 Fungal species in the Northwest Forest Plan. USDA, Forest Service Pacific Northwest Research Station, Portland, OR. GTR PNW-GTR-476. & Marr, C.D. and Stuntz, D.E. 1973. Ramaria of Western Washington. Biblio. Mycol. 38:1-232. Petersen, R.H. 1988. Contribution toward a monograph of Ramaria. VII. New Taxa and Miscellany. Mycologia 80:223-234

## New Sources