## **California Status Factors**

Elcode NFSM000164

**Gname** RAMARIA VERLOTENSIS

Gcomname

#### **Number of Occurrences**

A = 1 - 5

Comments There are 3 records from Coastal forests in northern CA.

## **Number of Occurrences with Good Viability**

B = Very few (1-3) occurrences with good viability

Comments There is a population in Jedediah Smith State Park that is protected.

## **Population Size**

A = 1-50 individuals

Comments

## **Range Extent**

D = 1,000-5,000 km2 (about 400-2,000 square miles)

Comments Known from Coastal forest in northern CA (3 locations)

## **Area of Occupancy**

A = <0.4 km2 (less than about 100 acres)

LA = <4 km (less than about 2.5 miles)

Comments

## Long-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Long-term trend in population, range, area occupied, or number or condition of occurrences unknown

Comments

# Short-term Trend in Population Size, Extent of Occurrence, Area of Occupancy, and/or Number or Condition of Occurrences

U = Unknown. Short-term trend in population, range, area occupied, and number and condition of occurrences unknown.

Comments

#### **Threats**

C = Substantial, non-imminent threat. Threat is moderate to severe but not imminent (> 10 years) for most of the population, occurrences, or area.

Scope High Severity High Immediacy Unknown

Comments This is a mycorrhizal species that depends on late successional forests of Douglas fir and western Hemlock

## **Number of Appropriately Protected and Managed Occurrences**

B = Few (1-3) occurrences appropriately protected and managed

Comments There is a population in Jedediah Smith State Park that is protected

## **Intrinsic Vulnerability**

A = Highly Vulnerable. Species is slow to mature, reproduces infrequently, and/or has low fecundity such that populations are very slow (> 20 years or 5 generations) to recover from decreases in abundance; or species has low dispersal capability such that extirpated populations are unlikely to become reestablished through natural recolonization (unaided by humans). Ecological community occurrences are highly susceptible to changes in composition and structure that rarely if ever are reversed through natural processes even over substantial time periods (> 100 years).

Comments This is a mycorrhizal species suceptible to logging activities

## **Environmental Specificity**

B = Narrow. Specialist or community with key requirements common.

Comments Mycorrhizal species

#### Other Considerations

Collections from CA may require reexamination to confirm their identity. If CA collections were misidentified then this species should be considered as G1. More studies are needed to determine its rarity or abundance.

Edition 11/11/2002 Edauthor Efren Cazares

**Grank** S2 **Grank Date** 11/11/2002

#### Greasons

There is only one out of 3 occurrences ia a protected site.

#### **BCD Sources**

Castellano, M.A., J.E. Smith, T. O'Dell, E. Cazares and S. Nugent. 1999. Handbook to Strategy 1 Fungal species in the Northwest Forest Plan. USDA, Forest Service Pacific Northwest Research Station, Portland, OR. GTR PNW-GTR-476. & Marr, C.D. and Stuntz, D.E. 1973. Ramaria of Western Washington. Biblio. Mycol. 38:1-232.

### **New Sources**